Check List the journal of biodiversity data

NOTES ON GEOGRAPHIC DISTRIBUTION

Check List 19 (4): 527–532
https://doi.org/10.15560/19.4.527

PENSOFT

New records of Largemouth Black Bass, *Micropterus salmoides* (Lacépède, 1802) (Pisces, Centrarchidae), in Bulgaria

Kiril Valkanov[1], Nikolay Kolev[2], Teodora Koynova[2*], Nikolay Natchev[2,3]

1 Department of Animal and Human Physiology, Faculty of Biology, Sofia University "St. Kliment Ohridski", Sofia, Bulgaria • kirilvalkanov55@gmail.com

2 Department of Biology, University of Shumen, Shumen, Bulgaria • NK: nd.kolev@hotmail.com • TK: t.koynova@shu.bg https://orcid.org/0000-0001-9044-6708 • NN: natchev@shu.bg https://orcid.org/0000-0002-0416-8738

3 Department of Evolutionary Zoology, University of Vienna, Vienna, Austria

* Corresponding author

Abstract. In the present study we report on new localities of Largemouth Black Bass, *Micropterus salmoides* (Lacépède, 1802), detected on the territory of Bulgaria. This invasive predator was observed for the first time in 2018 in a water basin in the capital city of Sofia city (western Bulgaria) and the following monitoring revealed that seven water bodies in the capital were inhabited by adults, subadults, and juvenile specimens. In 2021 and 2022, this species was captured in water basins in the vicinity of lake Mandra near the city of Burgas (south-eastren Bulgaria). Currently, these sites represent the easternmost localities within the distribution of *M. salmoides* in the Balkan peninsula and perhaps in continental Europe.

Keywords. Citizen science, invasive vertebrates, limnology, monitoring, predator, urban ecology

Academic editor: Arturo Angulo
Received 5 April 2023, accepted 23 June 2023, published 25 July 2023

Valkanov K, Kolev N, Koynova T, Natchev N (2023) New records of Largemouth Black Bass, *Micropterus salmoides* (Lacépède, 1802) (Pisces, Centrarchidae), in Bulgaria. Check List 19 (4): 527–532. https://doi.org/10.15560/19.4.527

Introduction

The Largemouth Black Bass, *Micropterus salmoides* (Lacépède, 1802) is a carnivorous perciform fish in the family Centrarchidae (sunfishes), which is native to Canada, USA, and Mexico in North America (De Vris et al. 2015). The species possesses a high potential for biological invasion and was included among the "100 most dangerous invasive species" list (Lowe et al. 2000). Since the 19th century, the fish was introduced as a commercial species in Africa, Asia, South America, and Europe (Pereira and Vitule 2019). *Micropterus salmoides* became a rather important fish food among the Asian human communities (Rojo-Cebreros et al. 2022) and currently China produces about 99% of the world's harvest of that aquaculture—about 0.7 million tonnes (Jin et al. 2022; Feng et al. 2023). In south-western Europe, *M. salmoides* was introduced as a game aquatic predator, which is predominantly of interest for sport fishing (Carpio et al. 2019), but also as a food source (Hrytsyniak and Guschin 2020). *Micropterus salmoides* were first released in France in 1877 (Drake 2009). Currently, the fish is distributed in Spain, Portugal, France, Italy, and other Mediterranean countries (Kottelat and Freyhof 2007; Carpio et al. 2019). On the Iberian Peninsula, the invasive potential of *M. salmoides* was identified as "high" (Almeida et al. 2013); however, in some parts of Europe (e.g., Ukraine), the species did not form stable populations, despite that it was imported more than a century ago (Kvach and Kutsokon 2017; Hrytsyniak and Guschin 2020).

On the Balkan peninsula, *M. salmoides* was introduced in 1892 in Slovenia (Seebens et al. 2017). In the last decades, this invasive fish was detected in Croatia (Dulčić et al. 2017), Bosna and Herzegovina (Tutman et al. 2017), Serbia (Seebens et al. 2017), and Romania (Stănescu et al. 2020; Iftime and Iftime 2021). This species was recently reported in Bulgaria—in the waters of the Struma River catchment area, but also northwestern from Sofia (Uzunova et al. 2019). In the present study,

©The authors. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

we provide new data on the presence of *M. salmoides* in the territory of Bulgaria and discuss the possible impact of this invasive species on the local population of fish and fish-based trophic webs.

Methods

Our investigation started in the spring of 2018. We performed monitoring of permanent water basins in the region of Sofia city in irregular intervals of about 12 days. We studied a total of 15 water bodies and performed a visual inspection for the presence of *M. salmoides*. In several cases, we were able to approach the fishes at a close distance and attempted to capture specimens with a handheld-fish nets (Fig. 1A). In late January 2023, we received information from local hobby-naturalists and fishers about the presence of *M. salmoides* in the region of Burgas city (Fig. 1B). The provided photographs were investigated by professional ichthyologists (KV, NK) and the taxonomy of the detected fishes was confirmed. The authors KV, NK, and NN identified the geographic position of the water bodies and visited the basins at the end of March 2023 (Fig. 2B, C). The water basins were coded as "1" and "2", and their position was matched to the database of the NATURA 2000 nets in Bulgaria (Fig. 2). The photographs by the authors were taken with a Panasonic Lumix DMC-FZ200 camera (Panasonic Corpo-ration, Osaka, Japan). All registrations were protocolled and georeferenced with the use of a handheld GPS System Garmin Etrex 30 GPS receiver (Garmin International Inc., Kansas, USA). The datum used for the recording of the geographical coordinates was WGS84. The exact coordinates of the already-published observations of *M. salmoides* in western Bulgaria (Fig. 2A) were provided by Uzunova et al. (2019). The updated map of the geographic distribution of *M. salmoides* was created using QGIS (QGIS Development Team 2021), and additional data were based on the dynamic map of NATURA 2000 PS in Bulgaria.

Results

Order Perciformes
Family Centrarchidae
Genus *Micropterus*

***Micropterus salmoides* (Lacépède, 1802)**

New records. BULGARIA – **Sofia** • 42.6592°N, 023.3076°E, 602 m a.s.l.; 19.VI.2020; KV captured 1 individual (Fig. 1A) within a water basin in South park • 42.6894°N, 023.4212°E, 509 m a.s.l. • 42.6893°N, 023.4202°E, 509 m a.s.l; • 42.6659°N, 023.4018°E, 544 a.s.l. • 42.6643°N, 023.4004°E, 544 m a.s.l. • 42.6637°N,

Figure 1. Largemouth black bass (*Micropterus salmoides*) in Bulgaria. **A.** Specimen of *M. salmoides* captured from a water basin in Sofia city. **B.** Two specimens captured at waterbody "1" by Hristo Valyarov on 10 September 2022.



Figure 2. Current distribution of the Largemouth black bass in Bulgaria. **A.** Map of the general distribution: black circles indicate the detected habitats of Uzunova et al. (2019), red dots indicate the newly registered habitats. **B.** Position of the two water basins indicated as "1" and "2" (in red) in the vicinity of Mandra lake. The territory of lake is protected from the of the NATURA 2000 network; however, water bodies "1" and "2" are positioned outside the protected site. **C.** Photograph of water body "1".

023.3991°E, 544 m a.s.l. • 42.6622°N, 023.3082°E, 616 m a.s.l • 42.6644°N, 023.3042°E, 616 m a.s.l.; 22.IV.2018–18.IX.2021; KV obs.; adults, subadults, juveniles – **Burgas** • Debelt • 42.3492°N, 027.2943°E, 90 m a.s.l.; water body "1" (Fig. 2B); 22.X.2021, Denis Toshev captured 1 individual; 10.IX.2022, Hristo Valjarov captured 2 individuals (Fig. 1B); 42.3831°N, 027.3147°E, 10 m a.s.l; water body "2", 15.IX.2022, Hristo Valjarov captured 4 individuals.

Identification. This species is the only representative of the genus *Micropterus* in Bulgaria. The only other centrarchid species that inhabits Bulgaria is the invasive Pumpkinseed Fish, *Lepomis gibbosus* (Linnaeus, 1758); however, it differs drastically in form and coloration from *M. salmoides* (Davis and Lock 2007). The fishes from the family Percidae, which inhabit the freshwaters of Bulgaria differ from *M. salmoides* in the coloration, the form of the body, the form of the fins, the body proportions, the position of the mouth, and the form and position of the eyes; for characteristics of these species, see Kottelat and Freyhof (2007) and van der Walt et al. (2019).

Discussion

According to Clavero and Garcia-Berthou (2005), biological invasions are one of the main factors or even the main cause for the extinction of animal species. *Micropterus salmoides* inflict crucial negative effects on the ecosystems, where it was introduced (Azuma 2001). Experimental data indicate that *M. salmoides* may radically impact species richness (67.86%) and biomass (up to 83.29%) of other fishes (Han et al. 2016). In the European freshwater basins, potential predators praying on *M. salmoides* can be *Esox lucius* Linnaeus, 1758 and some percids (Drake 2009). However, due to various factors, these predators cannot diminish the population of *M. salmoides* to critical levels. In many of the invaded waterbodies, *M. salmoides* becomes a top predator and, as such, it impacts by the "top-down" effect of various connections within the trophic webs (Han et al. 2016). The bass is specialized in the pursuit hunting of fish (Luger et al. 2020), and the levels of piscivory in this predator are among the highest recorded worldwide (Jang et al. 2006). The potential of this invasive predator to diminish dramatically the population of the local fish species is well documented and investigated (Han et al. 2016). Another rather alarming aspect is that *M. salmoides* is unable to "top-down" control the populations of other invaders, such as the sunfishes of the genus *Lepomis*. When these centrarchids coexist together, they prevail over the native fish species and create a great imbalance in the ichthyofauna, i.e., the sunfishes may appear in great numbers and totally dominate over the other fish species (Hossain et al. 2013). Another topic, that must be clearly understood is whether the European waters are invaded only by *M. salmoides*, or if Spotted Bass, *M. punctulatus* (Rafinesque, 1819) is also present (see van der Walt et al. 2019). Both species are closely related, externally look rather similar, and produce hybrids (Godbout et al. 2009). In eastern Europe, *M. salmoides* was detected rather sporadically. The species was recorded from the Volyne region in Ukraine (Kvach and Kutsokon 2017; Hrytsyniak and Guschin 2020) and in the region of Bucharest and two northern districts in Romania (Iftime and Iftime 2021). In this contribution, we report the presence of *M. salmoides* at the Black Sea coast in southeastern Bulgaria, being, to our best knowledge, the easternmost point of the range of that invasive species on the Balkan peninsula, or even in continental Europe.

The presence of *M. salmoides* at the very proximity to the Protected Site BG 0000271 Mandra lake is rather alarming; water basin "1" is located at about 4000 m and water basin "2" is less than 100 m from its boarder (Fig. 2B). Lake Mandra is a part of the so-called "Burgas lakes", or "Burgas wetlands", which are of great ecological importance. The Burgas lakes complex is highly sensitive and is recognized as Ramsar sites and is also a part of the Natura 2000 Protected Site network. It is important as permanent or transient habitat for a great variety of bird species, including many piscivorous species (Solakov 2010). Burgas wetlands are subject to intensive sport fishing, and the tendency in the last years is that the fishers practice the so called "catch and release" method. We propose that *M. salmoides* may spread in the system of lakes around Burgas city rather rapidly in the future. The Burgas wetlands are characterized by a strong variation in the water salinity—it ranges between 0.01 and 16.1‰ (Rabadjieva et al. 2019). While most freshwater fishes have a low tolerance toward increased conductivity of the waters they inhabit (Peterson and Meador 1994), *M. salmoides* is rather plastic concerning that factor (Susanto and Peterson 1996). Sun et al. (2021) explained the ability of this species to adapt to increased ionic pressure by the activation of a complex of genes, which switches a particular molecular mechanism for adaptation during the transition from fresh to brackish water. *Micropterus salmoides* is able to tolerate even seawater for at least some time (for an overview see Sun et al. 2021). The ability to inhabit waters with increased conductivity may allow for *M. salmoides* to invade large sections of the Burgas wetlands. The pressure from invasive carnivorous species may lead to drastic change in the behavior of highly protected species, such was the case with Lake "Srebarna" (Koshev et al. 2020). Having in mind the anthropogenic selective pressure and the genetic plasticity of *M. salmoides* (see Sun et al. 2023), we propose that *M. salmoides* may represent a severe threat for the fauna of the Burgas wetlands in the near future. Future field investigations using direct capture of the species and detection of its presence on the basis of environmental DNA (see Laramie et al. 2015) will provide detailed information on the spread of the invasive predator within the territory of Bulgaria. Urgent action plans for the elimination of *M. salmoides* must be issued and all good practices concerning its eradication must be applied immediately.

Acknowledgements

This work was supported by the Bulgarian Ministry of Education and Science under grant no. RD-08-80/09.02.2022 and no. RD-08-113/20.02.2023.

Author Contributions

Conceptualization: KV, NK, TK, NN. Data curation: NN. Funding acquisition: KV, NK, TK, NN. Investigation: KV, NK, NN. Methodology: KV, NN. Resources: KV, NK, NN. Visualization: TK, NN. Writing – original draft: KV, NN. Writing – review and editing: KV, NK, TK, NN.

References

Almeida D, Ribeiro F, Leunda P, Vilizzi L, Copp G (2013) Effectiveness of FISK, an invasiveness screening tool for non-native freshwater fishes, to perform risk identification assessments in the Iberian Peninsula. Risk Analysis 33 (8): 1404–1413. https://doi.org/10.1111/risa.12050

Azuma M (2001) Ecological disturbance by two alien fish species to the indigenous community of a small lake in western Japan, Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen 27 (7): 3794–3797. https://doi.org/10.1080/03680770.1998.11901693

Carpio AJ, De Miguel RJ, Oteros J, Hillstroem L, Tortosa F (2019) Angling as a source of non-native freshwater fish: a European review. Biological Invasions 21: 3233–3248. https://doi.org/10.1007/s10530-019-02042-5

Clavero M, Garcia-Berthou E (2005) Invasive species are a leading cause of animal extinctions. Trends in Ecology and Evolution 20 (3): 110. https://doi.org/10.1016/j.tree.2005.01.003

Davis JT, Lock JT (2007) Largemouth bass: biology and life history. Oklahoma Cooperative Extension Service.

De Vris DR, Wright RA, Glover DC, Farmer TM, Lowe MR, Norris AJ, Peer AC (2015) Largemouth bass in coastal estuaries: a comprehensive study from the Mobile–Tensaw River Delta, Alabama. In: Black Bass Diversity, Multidisciplinary Science for Conservation. American Fisheries Society Symposium 82: 297–310.

Dulčić J, Dragičević B, Ugarković P, Tutman P (2017) The largemouth black bass (*Micropterus salmoides*): first record in the Neretva River delta, Adriatic drainage system of Croatia. Cybium 41 (1): 77–78.

Drake A (2009) *Micropterus salmoides* (Largemouth Bass). CABI Compendium. https://doi.org/10.1079/cabicompendium.74846

Feng X, Yang H, Tu X, Wang Z Gu Z (2023) First report of Ambiphrya species (Ciliophora, Peritrichia) infestation in farmed *Micropterus salmoides* and *Ictalurus punctatus* in China. Aquaculture 563 (2): 738968. https://doi.org/10.1016/j.aquaculture.2022.738968

Godbout JD, Aday DD, Rice JA, Bangs MR, Quattro JM (2009) Morphological models for identifying largemouth Bass, Spotted Bass, and Largemouth Bass × Spotted Bass hybrids. North American Journal of Fisheries Management 29 (5): 1425–1437. https://doi.org/10.1577/m08-253.1

Han JH, Paek WK, An KG (2016) Exotic species, *Micropterus salmoides*, as a key bioindicator influencing the reservoir health and fish community structure. Journal of Asia-Pacific Biodiversity 9 (4): 403–411. https://doi.org/10.1016/j.japb.2016.10.002

Hrytsyniak I, Guschin V (2020) Aspects and Development of the Largemouth Bass (*Micropterus Salmoides*) Aquaculture in Ukraine. Annali d'Italia 13 (1): 3–7.

Hossain MM, Perhar G, Arhonditsis GB, Matsuishi T, Goto A, Azuma M (2013) Examination of the effects of largemouth bass (*Micropterus salmoides*) and bluegill (*Lepomis macrochirus*) on the ecosystem attributes of lake Kawahara-oike, Nagasaki, Japan. Ecological Informatics 18: 149–161. https://doi.org/10.1016/j.ecoinf.2013.07.005

Iftime A, Iftime O (2021) Alien fish, amphibian and reptile species in Romania and their invasive status: a review with new data. Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 64 (1): 131–186. https://doi.org/10.3897/travaux.64.e67558

Jin Y, Bergmann SM, Mai Q, Yang Y, Liu W, Sun D, Chen Y, Yu Y, Liu Y, Cai W, Dong H, Li H, Yu H, Wu Y, Lai M, Zeng W (2022) Simultaneous isolation and identification of largemouth bass virus and rhabdovirus from moribund largemouth bass (*Micropterus salmoides*). Viruses 14 (8): 1643. https://doi.org/10.3390/v14081643

Jang MH, Joo GJ, Lucas MC (2006) Diet of introduced largemouth bass in Korean rivers and potential interactions with native fishes. Ecology of Freshwater Fish 15 (3): 315–320. https://doi.org/10.1111/j.1600-0633.2006.00161.x

Koshev YS, Petrov MM, Nedyalkov NP, Raykov IA (2020) Invasive raccoon dog depredation on nests can have strong negative impact on the Dalmatian pelican's breeding population in Bulgaria. European Journal of Wildlife Research 66 (5): 85. https://doi.org/10.1007/s10344-020-01423-9

Kottelat M, Freyhof J (2007) Handbook of European Freshwater Fishes. Publications Kottelat, Cornol, Switzerland, 646 pp.

Kvach Y, Kutsokon Y (2017) The non-indigenous fishes in the fauna of Ukraine: a potentia ad actum. BioInvasions Records 6 (3): 269–279.

Laramie M, Pilliod D, Goldberg C, Strickler K (2015) Environmental DNA sampling protocol—filtering water to capture DNA from aquatic organisms. Techniques and Methods, 2–A13. United States Geological Survey, Reston, VA, USA. https://doi.org/10.3133/tm2A1

Lowe S, Browne M, Boudjelas S, De Poorter M (2000) 100 of the world's worst invasive alien species a selection from the Global Invasive Species Database. Aliens 12: 12.

Luger AM, South J, Alexander ME, Ellender BR, Weyl OLF, Nagelkerke LAJ (2020) Ecomorphology of largemouth bass relative to a native trophic analogue explains its high invasive impact. Biological Invasions 22 (7): 2223–2233. https://doi.org/10.1007/s10530-020-02252-2

Pereira FW, Vitule JRS (2019) The largemouth bass *Micropterus salmoides* (Lacepède, 1802): impacts of a powerful freshwater fish predator outside of its native range. Reviews in Fish Biology and Fisheries 29: 639–652. https://doi.org/10.1007/s11160-019-09570-2

Peterson M, Meador M (1994) Effects of salinity on freshwater fishes in coastal plain drainages in the southeastern US. Reviews in Fisheries Science 2 (2): 95–121. https://doi.org/10.1080/10641269409388554

Rabadjieva D, Kovacheva A, Tepavitcharova S, Gyuzelev G, Kornilev YV, Vasilev V (2018) Modeling of chemical species and precipitation processes in waters of the Protected Site Poda, Burgas, Bulgaria. Journal of International Scientific Publications, Ecology & Safety 12: 96–108.

Rojo-Cebreros A, Quintero H, Romano N, Heikes D Sanchez-Ramirez A (2022) Challenges in Largemouth Bass juvenile production. World Aquaculture 35: 34–39.

Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, Winter M, Arianoutsou M, Bacher S, Blasius B, Brundu G, Capinha C, Celesti-Grapow L, Dawson W, Dullinger S, Fuentes N, Jäger H, Kartesz J, Kenis M, Kreft H, Kühn I, Lenzner B, Liebhold A, Mosena A, Moser D, Nishino M, Pearman D, Pergl J, Rabitsch W, Rojas-Sandoval J, Roques A, Rorke S, Rossinelli S, Roy HE, Scalera R, Schindler S, Štajerová K, Tokarska-Guzik B, van Kleunen M, Walker K, Weigelt P, Yamanaka T, Essl F (2017) No saturation in the accumulation of alien species worldwide. Nature

Communications 8: 14435. https://doi.org/10.1038/ncomms14435

Solakov D (2010) Wetlands of international importance for Bulgaria. NEAR Curriculum in Natural Environmental Science. Terre et Environnement 88: 129–135.

Stănescu F, Rozylowicz L, Tudor M, Cogălniceanu D (2020) Alien vertebrates in Romania—a review. Acta Zoologica Bulgarica 72 (4): 583–595.

Sun C, Li J, Dong J, Niu Y, Hu J, Lian J, Li W, Li J, Tian Y, Shi Q, Ye X (2021) Chromosome-level genome assembly for the largemouth bass *Micropterus salmoides* provides insights into adaptation to fresh and brackish water. Molecular Ecology Resources 21 (1): 301–315. https://doi.org/10.1111/1755-0998.13256

Sun C, Zhang XH, Dong JJ, You XX, Tian YY, Gao FY, Zhang HT, Shi Q, Ye X (2023) Whole-genome resequencing reveals recent signatures of selection in five populations of largemouth bass (*Micropterus salmoides*). Zoological Research 44 (1): 78–89. https://doi.org/10.24272/j.issn.2095-8137.2022.274

Susanto GN, Peterson MS (1996) Survival, osmoregulation and oxygen consumption of YOY coastal largemouth bass, *Micropterus salmoides* (Lacepede) exposed to saline media. Hydrobiologia 323: 119–127. https://doi.org/10.1007/bf00017589

Tutman P, Hamzić A, Džano A, Dulčić J (2017) The first record of non-native largemouth black bass, *Micropterus salmoides* (Actinopterygii: Perciformes: Centrarchidae), in Bosnia and Herzegovina. Acta Ichthyologica et Piscatoria 47 (3): 317–320. https://doi.org/10.3750/aiep/02187

Uzunova E, Studenkov S, Dashinov D (2019) First records of largemouth bass *Micropterus salmoides* (Lacépède, 1802) from Bulgaria (Balkan Peninsula). BioInvasions Records 8 (2): 427–436. https://doi.org/10.3391/bir.2019.8.2.25

van der Walt JA, Marr SM, Wheeler MJ, Impson ND, Garrow C, Weyl OL (2019) Successful mechanical eradication of spotted bass (*Micropterus punctulatus* (Rafinesque, 1819)) from a South African river. Aquatic Conservation: Marine and Freshwater Ecosystems 29 (2): 303–311. https://doi.org/10.1002/aqc.3035